

LIGHT QUARK MASSES FROM QUENCHED LATTICE QCD SIMULATIONS WITH DOMAIN WALL QUARKS*

MATTHEW WINGATE

*RIKEN BNL Research Center, Brookhaven National Laboratory
Upton, NY 11973, USA*

Values for the strange quark mass and average up/down mass have been obtained from quenched lattice QCD simulations using the domain wall fermion action. This discretization preserves the properties of flavor and chiral symmetry at nonzero lattice spacing. Results are shown for two values of the lattice spacing. The mass renormalization constant is computed nonperturbatively.

The light quark masses are fundamental parameters of the Standard Model and, as such, must be determined more precisely in order to increase its predictive power. Lattice QCD can play an important role in computing the quark masses, as it already has with the strong coupling constant. Recently it was shown that the new domain wall fermion discretization can be applied to this endeavor.¹ This talk reports on the progress of bringing these calculations to the present level of other lattice quark mass calculations.^{2,3}

The RBC Collaboration has recently reported quenched lattice QCD results for light hadron masses and matrix elements from a series of simulations.⁴ We used domain wall quarks, which approximately maintain continuum chiral symmetry by introducing a fifth dimension of size L_s .⁵ The nearly chiral surface states at the boundaries mix within the bulk of the fifth dimension. This breaking effect can be parameterized for low energy matrix elements in terms of a residual quark mass m_{res} . A tunable quark mass m_f is introduced to the action which explicitly couples the surface states, thus one studies operators containing quarks with effective masses equal to $m_f + m_{\text{res}}$.

*Talk given at DPF 2000, Columbus, OH, USA; work done in collaboration with T. Blum, P. Chen, N. Christ, C. Cristian, C. Dawson, G. Fleming, A. Kaehler, X. Liao, G. Liu, C. Malureanu, R. Mawhinney, S. Ohta, G. Siegert, A. Soni, C. Sui, P. Vranas, L. Wu, and Y. Zhestkov (RIKEN/BNL/CU Collaboration)




Figure 1: Axial WT identity quark mass (lattice units and regularization) vs. m_f . Dashed lines indicate where the physical K or K^* mass is obtained.




Figure 2: Z_A/Z_P in RI/MOM scheme at 2 GeV, plotted vs. input momentum squared. Line and asterisk indicate linear extrapolation to $(ap)^2 = 0$ to remove any cutoff effects.

Here we report results from simulations using two values of the gauge coupling parameter $\beta \equiv 6/g_0^2$. The lattice spacing is set through M_ρ . The m_f corresponding to the physical strange sector, $m_f^{(s)}$, is set by requiring either the pseudoscalar meson mass to equal M_K or the vector mass to equal M_{K^*} . It is a common feature of quenched simulations that $m_f^{(s)}$ from the K is smaller than from K^* , so results are quoted for both inputs. The simulations here are done with degenerate quarks, so $m_f^{(s)} \approx m_s/2$. Simulations with nondegenerate quarks show SU(3) flavor symmetry is preserved within errors.

In Fig. 1 we show the quark mass obtained through the axial Ward-Takahashi identity (AWTI) by directly computing matrix elements of the local axial current and pseudoscalar density. The renormalization factor for this definition is Z_A/Z_P . We compute this nonperturbatively in the RI/MOM scheme⁶ for several momenta; the results are plotted in Fig. 2.⁷

The quark mass may also be defined through the vector Ward-Takahashi identity (VWTI). The renormalization constant for this definition is Z_V/Z_S .

Table 1. Strange quark mass in MeV ($\overline{\text{MS}}, 2 \text{ GeV}$) $\pm$ stat. $\pm$ sys. Scale set through M_ρ .

β	K input		K^* input	
	VWTI	AWTI	VWTI	AWTI
5.85	–	100(5)(20)	–	138(7)(28)
6.00	110(2)(22)	105(6)(21)	132(2)(26)	127(6)(25)

Rather than directly computing matrix elements, one can use the difference $2(m_f^{(s)} - m_{\text{res}}) = m_s + m_{u,d}$ (we assume⁸ $m_{u,d} = m_s/24.4$). Since the vector current matrix element is an intermediate step, we are free to choose the conserved one, for which $Z_V = 1$. The calculation of Z_S is harder since the wave function renormalization Z_q , which cancels in the ratio Z_A/Z_P , now needs to be computed. So far, this has been done precisely only for $\beta = 6.00$.

Table 1 summarizes our results. The conversion to the $\overline{\text{MS}}$ scheme was done perturbatively at two loops. At 2 GeV the three loop term is a sizable correction,⁹ which we include as a 5% systematic error. The only way to reduce this uncertainty is to do the matching, and thus the whole simulation, with a scale closer to 3 GeV. The $\approx 10\%$ systematic uncertainty from the lattice spacing is not included, but should be if one uses these values for phenomenology.

The results presented here are updated from last year.¹⁰ The strange quark mass given then was taken using the VWTI, and a large uncertainty was assigned due to the difference between $-m_{\text{res}}$ and the m_f where $M_\pi^2 \rightarrow 0$ linearly. Now that we have seen how topological zero modes and quenching effects distort the linear extrapolation to the chiral limit,⁴ this uncertainty is removed.

These are early steps in bringing quenched domain wall calculations to the level of those with other lattice actions. Simulations with finer lattice spacings are necessary to take the continuum limit, and then an important check of systematic effects between different discretizations can be made.

Acknowledgments

Simulations done on the RBRC and CU QCDSP's and the NERSC Cray T3E.

References

1. T. Blum, A. Soni and M. Wingate, Phys. Rev. **D60** (1999) 114507.
2. R. Gupta, these proceedings.
3. V. Lubicz, plenary talk at Lattice 2000.
4. T. Blum *et al.*, hep-lat/0007038.
5. For a review and references see P. Vranas, plenary talk at Lattice 2000.
6. G. Martinelli *et al.*, Nucl. Phys. **B445** (1995) 81.
7. T. Blum *et al.*, in preparation; see also C. Dawson, talk at Lattice 2000.
8. H. Leutwyler, hep-ph/9609467.
9. K. G. Chetyrkin and A. Rétey, Nucl. Phys. **B583** (2000) 3.
10. M. Wingate, Nucl. Phys. Proc. Suppl. **83** (2000) 221 [hep-lat/9909101].